



University of Tennessee, Knoxville

TRACE: Tennessee Research and Creative Exchange

Insects, Pests, Plant Diseases and Weeds

UT Extension Publications

5-14-2013

W 289-O IPM QuickFacts Series: Apple Scab

Amy Fulcher

Heather Bowers

Follow this and additional works at: https://trace.tennessee.edu/utk_agexdise

 Part of the [Agricultural Education Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

"W 289-O IPM QuickFacts Series: Apple Scab," Amy Fulcher and Heather Bowers,
W 289-O
, https://trace.tennessee.edu/utk_agexdise/76

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Ornamental insects, Pests & Diseases is brought to you for free and open access by the UT Extension Publications at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Insects, Pests, Plant Diseases and Weeds by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Apple Scab *Venturia inaequalis*



Introduction

Apple scab is a serious disease of apples and crabapples. Repeated spring and summer defoliation results in fewer flowers the next spring and overall weakening of the tree. For apple trees, apple scab can greatly reduce yield through repeated defoliation and unsalable fruit.

Host Plants

- Apple
- Crabapple
- Hawthorn
- Mountain ash

Pathogen and Disease Cycle

Apple scab is caused by the fungus *Venturia inaequalis*. There are two stages of this disease cycle: the primary infection and the secondary infection.

Primary infection

The fungus overwinters on fallen infected leaves. In the spring during the rainy season, fungal spores are released from the infected leaves that remain from the previous year and are carried by the wind to newly emerging leaves and fruits of healthy trees.

Secondary infection

Once primary infection occurs, secondary spores (conidia) are formed, allowing a tree to continually reinfect itself or neighboring trees as long as environmental conditions are favorable. The disease progresses rapidly and causes severe infections if plants remain continuously wet for at least a six-hour period with temperatures around 65 to 75 degrees F. The warmer the temperature, the less time the leaf must remain wet continuously for infection to occur.

Monitoring

Watch for initial leaf symptoms during the pre-bloom period. Examine the upper and lower leaf surfaces, in particular leaf veins, as this area will retain moisture longer, meeting the moisture requirements for spore germination faster than other areas of the leaf. Be sure to note durations of wet periods and average temperatures when fungal spores are produced in early spring and compare to charts for spore germination (see UK PPA-24) to anticipate infection periods.



Symptoms



Leaves

Infection first appears as olive green spots with fuzzy margins on both sides of infected leaves. As the disease progresses, the spots develop a more prominent outline and become black and velvety. Leaves will ultimately turn yellow and drop prematurely.

Fruit

Fruit infection symptoms are similar to those of leaves. Lesions on fruit turn dark brown to black. As the fruit develops, lesions will have a “scabby” appearance and will crack as fruit grows larger. Severe infections while fruit is small can cause deformity from uneven growth.



Integrated Pest Management

CULTURAL CONTROL

Plant resistant apple cultivars such as Enterprise, Freedom, Gold Rush, Jonafree, Liberty, Macfree, Nova Easy, Prima, Priscilla, Redfree, Sir Prize and William’s Pride. Plant resistant crabapple cultivars (see J. Hartman publication PPA-24 below). Prune trees to allow better air penetration, which will help keep plants from staying moist continuously. Dispose of fallen leaves in autumn.

CHEMICAL CONTROL

Please refer to http://eppserver.ag.utk.edu/redbook/sections/trees_flowers.htm for the most up-to-date recommendations.

Resources

Photo credits: Steve Bost and Alan Windham, University of Tennessee

Biggs, A.R. and K.D. Hickey. Apple scab *Venturia inaequalis*. Kearneysville Tree Fruit Research and Education Center.

http://www.caf.wvu.edu/kearneysville/disease_descriptions/omapscab.html

Hartman, J. Apple scab. University of Kentucky Extension publication PPA-24. <http://www.ca.uky.edu/agc/pubs/ppa/ppa24/ppa24.htm>

Schroeder, D. and E. Marotte. 2003. Apple scab. University of Connecticut Integrated Pest Management Program.

<http://www.hort.uconn.edu/ipm/homegmd/htms/19apscab.htm>

Snover-Clift, K. and S. Jensen-Tracy. *Venturia inaequalis*. Bugwoodwiki. http://wiki.bugwood.org/Venturia_inaequalis

THE UNIVERSITY of TENNESSEE 
INSTITUTE of AGRICULTURE

Prepared by Heather Bowers, Intern, Department of Plant Sciences
and Dr. Amy Fulcher, Assistant Professor, Department of Plant Sciences

Publication funded by USDA Extension IPM Grant in partnership with University of Kentucky Integrated Pest Management Program.

The authors thank M. Halcomb, S. Bost, W. Russell and A. Windham for their careful review.

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.
UT Extension provides equal opportunities in programs and employment.